

Lattice uniformities on effect algebras

Anna Avallone and Paolo Vitolo

Dipartimento di Matematica, Università della Basilicata
Contrada Macchia Romana, 85100 Potenza, Italy
avallone@unibas.it, vitolo@unibas.it

Abstract

Let L be a lattice ordered effect algebra. We prove that the lattice uniformities on L which make uniformly continuous the operations $\ominus$ and $\oplus$ of L are uniquely determined by their system of neighbourhoods of 0 and form a distributive lattice. Moreover we prove that every such uniformity is generated by a family of weakly subadditive $[0, +\infty]$ -valued functions on L .

Keywords Effect algebras, lattice uniformities, submeasures.

Introduction

Effect algebras have been introduced by D. J. Foulis and M. K. Bennett in 1994 [8] for modelling unsharp measurement in a quantum mechanical system. They are a generalization of many structures which arise in quantum physics (see [7]) and in Mathematical Economics (see [11, 9]), in particular of orthomodular lattices in non-commutative measure theory and MV-algebras in fuzzy measure theory. After 1994, there have been a great number of papers concerning effect algebras (see [10] for a bibliography).

In this paper we study D-uniformities on a lattice ordered effect algebra L , i.e. lattice uniformities on L which make uniformly continuous the operations $\ominus$ and $\oplus$ of L .

The starting point of our paper is observing the key role played by D-uniformities in the study of modular measures on L (see [1, 2, 3]), since every modular measure on L generates a D-uniformity. Also of importance is the role played in the study of modular functions on orthomodular lattices (see [16]) and of measures on MV-algebras (see [6, 13]) by the lattice structure of filters which generate lattice uniformities making uniformly continuous the operations of these structures.

In the first part of the paper, we give a description of the filters which are systems of neighbourhoods of 0 in D-uniformities on L —called D-filters—and we prove that there exists an order isomorphism between the lattice of all D-uniformities on L and the lattice of all D-filters on L . In particular every D-uniformity is uniquely determined by its system of neighbourhoods of 0. As a consequence, we obtain that the lattice of all D-uniformities on L is distributive.

Our results extend similar results of [16] in orthomodular lattices (see also [5]) and of [6] and [13] in MV-algebras, and give as particular case the order isomorphism found in [4] between some lattice congruences and some lattice ideals.

In the second part of the paper, we apply the results of the first part to prove that every D-uniformity on L is generated by a family of weakly subadditive $[0, +\infty]$ -functions on L .

1 Preliminaries

An *effect algebra* [10] is a set E , with two distinguished elements 0 and 1, and a partially defined operation $\oplus$ such that for all $a, b, c \in E$:

- (E1) If $a \oplus b$ is defined, then $b \oplus a$ is defined and $a \oplus b = b \oplus a$.
- (E2) If $b \oplus c$ is defined and $a \oplus (b \oplus c)$ is defined, then $a \oplus b$ and $(a \oplus b) \oplus c$ are defined, and $a \oplus (b \oplus c) = (a \oplus b) \oplus c$.
- (E3) There exists a unique $a^\perp \in E$ such that $a \oplus a^\perp$ is defined and $a \oplus a^\perp = 1$.
- (E4) If $a \oplus 1$ is defined, then $a = 0$.

It is easily seen that $a \oplus 0$ is always defined and equals a . If $a \oplus b$ is defined, we say that a and b are *orthogonal* and write $a \perp b$.

In an effect algebra E another partially defined operation $\ominus$ can be defined by the following rule: $c \ominus a$ exists and equals b if and only if $a \oplus b$ exists and equals c . In particular, $a^\perp = 1 \ominus a$. Moreover, if $a \perp b$, then $a \oplus b = (a^\perp \ominus b)^\perp = (b^\perp \ominus a)^\perp$.

In an effect algebra E a partial ordering relation $\leq$ can be defined as follows: $a \leq c$ if and only if, for some $b \in E$, $a \oplus b$ exists and equals c . Hence $c \ominus a$ is defined if and only if $a \leq c$. Moreover $a \perp b$ if and only if $a \leq b^\perp$.

If $a \vee b$ and $a \wedge b$ exist for all $a, b \in E$, then we say that E is a *lattice ordered effect algebra* (otherwise called *D-lattice*). In this case, we define the *symmetric difference* of any two elements a and b in E as $a \triangle b = (a \vee b) \ominus (a \wedge b)$.

Throughout the paper, the symbol L will always denote a lattice ordered effect algebra. Let us recall that L is an *MV-algebra* if and only if $(a \vee b) \ominus b = a \ominus (a \wedge b)$ for all $a, b \in L$, while L is an *orthomodular lattice* if and only if $a^\perp \wedge a = 0$ for every $a \in L$.

We will make use of the following properties (for the proofs we refer to [10]).

1.1. Proposition. *For all $a, b, c \in L$ we have:*

- (i) *If $a \leq b$, then $b \ominus a \leq b$ and $b \ominus (b \ominus a) = a$.*
- (ii) *If $a \leq b \leq c$, then $c \ominus b \leq c \ominus a$ and $(c \ominus a) \ominus (c \ominus b) = b \ominus a$.*
- (iii) *If $a \leq b \leq c$, then $b \ominus a \leq c \ominus a$ and $(c \ominus a) \ominus (b \ominus a) = c \ominus b$.*
- (iv) *If $a \leq b^\perp$ and $a \oplus b \leq c$, then $c \ominus (a \oplus b) = (c \ominus a) \ominus b = (c \ominus b) \ominus a$.*
- (v) *If $a \leq b \leq c^\perp$, then $a \oplus b \leq b \oplus c$ and $(b \oplus c) \ominus (a \oplus c) = b \ominus a$.*
- (vi) *If $a \leq b \leq c$, then $a \oplus (c \ominus b) = c \ominus (b \ominus a)$.*
- (vii) *If $a \leq b^\perp \leq c^\perp$, then $a \oplus (b \ominus c) = (a \oplus b) \ominus c$.*
- (viii) *If $a \leq c$ and $b \leq c$, then $c \ominus (a \vee b) = (c \ominus a) \wedge (c \ominus b)$ and $c \ominus (a \wedge b) = (c \ominus a) \vee (c \ominus b)$.*
- (ix) *If $c \leq a$ and $c \leq b$, then $(a \wedge b) \ominus c = (a \ominus c) \wedge (b \ominus c)$ and $(a \vee b) \ominus c = (a \ominus c) \vee (b \ominus c)$.*
- (x) *If $a \leq c^\perp$ and $b \leq c^\perp$, then $(a \vee b) \oplus c = (a \oplus c) \vee (b \oplus c)$ and $(a \wedge b) \oplus c = (a \oplus c) \wedge (b \oplus c)$.*

Let $\mathcal{U}$ be a uniformity on L . We say that $\mathcal{U}$ is a *lattice uniformity* [14] if the operations $\vee$ and $\wedge$ are uniformly continuous with respect to $\mathcal{U}$.

A *D-uniformity* [1] is a lattice uniformity which makes the operations $\oplus$ and $\ominus$ uniformly continuous, too. The set of all D-uniformities on L will be denoted by $\mathcal{DU}(L)$. It is easy to see that $\mathcal{DU}(L)$ —ordered by inclusion—is a complete lattice, with the discrete uniformity and the trivial uniformity as greatest and smallest elements, respectively.

Given $U, V \subset L \times L$, we put

$$\begin{aligned} U \vee V &= \{ (a_1 \vee b_1, a_2 \vee b_2) : (a_1, a_2) \in U, (b_1, b_2) \in V \}, \\ U \wedge V &= \{ (a_1 \wedge b_1, a_2 \wedge b_2) : (a_1, a_2) \in U, (b_1, b_2) \in V \}, \\ U \ominus V &= \{ (a_1 \ominus b_1, a_2 \ominus b_2) : b_1 \leq a_1, b_2 \leq a_2, (a_1, a_2) \in U, (b_1, b_2) \in V \}. \end{aligned}$$

It is known (see [14]) that a uniformity $\mathcal{U}$ on L is a lattice uniformity if and only if for every $U \in \mathcal{U}$ there exists $V \in \mathcal{U}$ such that $V \vee \Delta \subset U$ and $V \wedge \Delta \subset U$, where $\Delta = \{ (a, a) : a \in L \}$.

Similarly, it has been shown in [1] that a lattice uniformity $\mathcal{U}$ on L is a D-uniformity if and only if for every $U \in \mathcal{U}$ there exists $V \in \mathcal{U}$ such that $V \ominus \Delta \subset U$ and $\Delta \ominus V \subset U$.

2 D-uniformities and D-filters

2.1. Definition. A filter $\mathcal{F}$ of subsets of a D-lattice L is called a *D-filter* if it satisfies the following:

- (F1) $\forall F \in \mathcal{F} \quad \exists F' \in \mathcal{F} : \quad \forall a, b \in F' \quad [a \perp b \implies a \oplus b \in F];$
- (F2) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall a \in G \quad \forall c \in L \quad (a \vee c) \ominus c \in F.$

The set of all D-filters on L will be denoted by $\mathcal{FND}(L)$.

Note that, by 1.1(ii), a filter $\mathcal{F}$ satisfies (F2) if and only if, for every $F \in \mathcal{F}$, there exists $G \in \mathcal{F}$ such that, for all $a \in G$ and all $c \in L$, one has $c \ominus (a^\perp \wedge c) \in F$.

We shall prove, in Theorem 2.4 below, that $\mathcal{FND}(L)$ is isomorphic to $\mathcal{DU}(L)$ and that $\mathcal{F}$ is a D-filter if and only if $\mathcal{F}$ is the system of neighbourhoods of 0 in a D-uniformity.

2.2. Lemma. *For every $a, b, c, d \in L$ such that $c \leq a$, $c \leq b$, $d \geq a$ and $d \geq b$ one has $(a \ominus c) \triangle (b \ominus c) = a \triangle b = (d \ominus a) \triangle (d \ominus b)$.*

PROOF. Indeed, applying 1.1(ix), and 1.1(iii), one gets $(a \ominus c) \triangle (b \ominus c) = ((a \ominus c) \vee (b \ominus c)) \ominus ((a \ominus c) \wedge (b \ominus c)) = ((a \vee b) \ominus c) \ominus ((a \wedge b) \ominus c) = (a \vee b) \ominus (a \wedge b) = a \triangle b$. Similarly, applying 1.1(viii), and 1.1(ii), one gets $(d \ominus a) \triangle (d \ominus b) = ((d \ominus a) \vee (d \ominus b)) \ominus ((d \ominus a) \wedge (d \ominus b)) = (d \ominus (a \wedge b)) \ominus (d \ominus (a \vee b)) = (a \vee b) \ominus (a \wedge b) = a \triangle b$. $\square$

2.3. Proposition. *A D-filter $\mathcal{F}$ on L has the following properties:*

- (i) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall a \in G \quad \forall b \in L \quad [b \leq a \implies b \in F]$;
- (ii) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall a, b \in G \quad a \vee b \in F$;
- (iii) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall x, y, z \in L \quad [x \triangle y \in G \implies (x \vee z) \triangle (y \vee z) \in F]$;
- (iv) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall x, y, z \in L \quad [x \triangle y \in G \implies (x \wedge z) \triangle (y \wedge z) \in F]$;
- (v) $\forall F \in \mathcal{F} \quad \exists G \in \mathcal{F} : \quad \forall x, y, z \in L \quad [x \triangle y \in G, y \triangle z \in G \implies x \triangle z \in F]$.

PROOF.

(i) Let $F \in \mathcal{F}$ and let $G \in \mathcal{F}$ such that (F2) is satisfied. Given any $a \in G$ and any $b \in L$ with $b \leq a$, put $c = a \ominus b$. Then $b = a \ominus (a \ominus b) = (a \vee (a \ominus b)) \ominus (a \ominus b) = (a \vee c) \ominus c \in F$.

(ii) Given $F \in \mathcal{F}$, let $F' \in \mathcal{F}$ satisfy (F1), and let $G \in \mathcal{F}$ satisfy (F2) with F' in place of F . If $a, b \in G$, then $(a \vee b) \ominus b \in F'$. Moreover $b \in F'$ by (i). Therefore $a \vee b = ((a \vee b) \ominus b) \oplus b \in F$.

(iii) Let $F \in \mathcal{F}$ and let $G \in \mathcal{F}$ such that (F2) is satisfied. Given x, y, z such that $x \triangle y \in G$, we put $a = x \triangle y$ and $c = ((x \vee z) \wedge (y \vee z)) \ominus (x \wedge y)$ and we show that $(x \vee z) \triangle (y \vee z) = (a \vee c) \ominus c$. First observe that $x \vee y \vee z = (x \vee y) \vee ((x \vee z) \wedge (y \vee z))$. Now, applying 1.1(x) and 1.1(v), we have:

$$\begin{aligned} (x \vee z) \triangle (y \vee z) &= (x \vee y \vee z) \ominus ((x \vee z) \wedge (y \vee z)) \\ &= (((x \vee y) \vee ((x \vee z) \wedge (y \vee z))) \ominus ((x \vee z) \wedge (y \vee z))) \\ &= (((x \triangle y) \oplus (x \wedge y)) \vee ((x \vee z) \wedge (y \vee z))) \ominus ((x \vee z) \wedge (y \vee z)) \\ &= ((a \oplus (x \wedge y)) \vee (c \oplus (x \wedge y))) \ominus (c \oplus (x \wedge y)) \\ &= ((a \vee c) \oplus (x \wedge y)) \ominus (c \oplus (x \wedge y)) = (a \vee c) \ominus c. \end{aligned}$$

(iv) Given $F \in \mathcal{F}$, take $G \in \mathcal{F}$ such that (iii) is satisfied, and let x, y, z such that $x \triangle y \in G$. By Lemma 2.2 we have $x^\perp \triangle y^\perp = x \triangle y$, and therefore $(x \wedge z) \triangle (y \wedge z) = (x^\perp \vee z^\perp)^\perp \triangle (y^\perp \vee z^\perp)^\perp = (x^\perp \vee z^\perp) \triangle (y^\perp \vee z^\perp) \in F$.

(v) Given $F \in \mathcal{F}$, let $F_1 \in \mathcal{F}$ satisfy (ii), let $F_2 \in \mathcal{F}$ satisfy (i) with F_1 in place of F , let $F_3 \in \mathcal{F}$ satisfy (F1) with F_2 in place of F and let $G \in \mathcal{F}$ satisfy (iii) with F_3 in place of F . If $a, b, c \in L$ are such that both $x \triangle y$ and $x \triangle z$ belong to G , then $a = (x \vee y \vee z) \ominus (y \vee z) = ((x \vee (x \vee z)) \triangle (y \vee (y \vee z))) \in F_3$ and $b = (y \vee z) \ominus z = (y \vee z) \triangle (z \vee z) \in F_3$ also. It follows that $(x \vee y \vee z) \ominus z = a \oplus b \in F_2$, so that $(x \vee z) \ominus z \in F_1$. Similarly one shows that $(x \vee z) \ominus x \in F_1$. Hence $x \triangle z = ((x \vee z) \ominus z) \vee ((x \vee z) \ominus x) \in F$. $\square$

2.4. Theorem.

- (a) *If $\mathcal{U}$ is a D-uniformity, then the filter $\mathcal{F}_{\mathcal{U}}$ of neighbourhoods of 0 in $\mathcal{U}$ is a D-filter.*
- (b) *Let $\mathcal{F}$ be a D-filter and, for each $F \in \mathcal{F}$, let $F^\triangle = \{(a, b) \in L \times L : a \triangle b \in F\}$. Then $\mathcal{B} = \{F^\triangle : F \in \mathcal{F}\}$ is a base for a D-uniformity whose filter of neighbourhoods of 0 is $\mathcal{F}$.*
- (c) *The mapping $\Psi : \mathcal{U} \mapsto \mathcal{F}_{\mathcal{U}}$ is an order-isomorphism of $\mathcal{DU}(L)$ onto $\mathcal{FND}(L)$ (both ordered by inclusion).*

PROOF.

(a) Since $\oplus$ is continuous at $(0,0)$, for every $F \in \mathcal{F}_U$ there exists $F' \in \mathcal{F}_U$ such that if $(a,b) \in F' \times F'$ and $a \perp b$, then $a \oplus b \in F$. This gives (F1). To prove (F2), let $F \in \mathcal{F}_U$ and let $U \in \mathcal{U}$ with $U(0) \subseteq F$. By uniform continuity of $\ominus$ and $\vee$, there exist $V_1, V_2 \in \mathcal{U}$ such that $V_1 \ominus \Delta \subset U$ and $V_2 \vee \Delta \subset V_1$. Now put $G = V_2(0)$, and consider any $a \in G$. Then $(0,a) \in V_2$, so that for every $c \in L$ we have $(c, a \vee c) \in V_2 \vee \Delta \subset V_1$ and hence $(0, (a \vee c) \ominus c) \in V_1 \ominus \Delta \subset U$ which means that $(a \vee c) \ominus c \in U(0) \subseteq F$.

(b) Clearly F^Δ is symmetric and $\Delta \subset F^\Delta$ for every $F \in \mathcal{F}$. Moreover, given $F_1, F_2 \in \mathcal{F}$, let $F_3 = \mathcal{F}_1 \cap F_2$. Then $F_3^\Delta = F_1^\Delta \cap F_2^\Delta$. Finally, if $F \in \mathcal{F}$ and $G \in \mathcal{F}$ satisfies 2.3(v), we have that $G^\Delta \circ G^\Delta \subseteq F^\Delta$. Therefore $\mathcal{B}$ is a base for a uniformity $\mathcal{U}$.

Now fix $U \in \mathcal{U}$. We show that there exists $V \in \mathcal{U}$ such that both $V \vee \Delta$ and $V \wedge \Delta$ are contained in U . Let $G \in \mathcal{F}$ satisfy 2.3(iii) and put $V = G^\Delta$. Given $(x,y) \in V \vee \Delta$, take $a, b, c \in L$ with $x = a \vee c$, $y = b \vee c$ and $(a,b) \in V$, that is $a \Delta b \in G$. By 2.3(iii), we have $x \Delta y = (a \vee c) \Delta (b \vee c) \in F$, that is $(x,y) \in F^\Delta$. We conclude that $V \vee \Delta \subset F^\Delta \subseteq U$. Since the same G also satisfies 2.3(iv) one sees in a similar way that $V \wedge \Delta \subset F^\Delta \subseteq U$ too. Next, we show that there exists $V \in \mathcal{U}$ such that both $V \ominus \Delta$ and $\Delta \ominus V$ are contained in U . Choose $F \in \mathcal{F}$ such that $F^\Delta \subseteq U$ and put $V = F^\Delta$. By Lemma 2.2, one has $F^\Delta \ominus \Delta = \{(a \ominus c, b \ominus c) : c \leq a, c \leq b, a \Delta b \in F\} = \{(a \ominus c, b \ominus c) : c \leq a, c \leq b, (a \ominus c) \Delta (b \ominus c) \in F\} = F^\Delta$ and similarly one sees that $\Delta \ominus F^\Delta = F^\Delta$. Hence $V \ominus \Delta \subset U$ and $\Delta \ominus V \subset U$.

It remains to prove that the filter of neighbourhoods of 0 in $\mathcal{U}$ coincides with $\mathcal{F}$. First observe that, given any $F \in \mathcal{F}$, we have

$$F^\Delta(0) = \{a \in L : (0,a) \in F^\Delta\} = \{a \in L : a \Delta 0 \in F\} = F \quad (2;1)$$

and therefore F is a neighbourhood of 0 in $\mathcal{U}$. Conversely, if G is a neighbourhood of 0 in $\mathcal{U}$, since $\mathcal{B}$ is a base for $\mathcal{U}$, there exists $F \in \mathcal{F}$ such that $F^\Delta(0) \subseteq G$. By (2;1), this means that $F \subset G$ and hence $G \in \mathcal{F}$, because $\mathcal{F}$ is a filter.

(c) It follows from (a) that Ψ maps $\mathcal{DU}(L)$ into $\mathcal{FND}(L)$. Now for any $\mathcal{F} \in \mathcal{FND}(L)$ let $\Phi(\mathcal{F})$ denote the D-uniformity constructed as in (b). Since $\Psi(\Phi(\mathcal{F})) = \mathcal{F}$, we have that Ψ is onto. Moreover if $\mathcal{F}_1, \mathcal{F}_2 \in \mathcal{DU}(L)$ and $\mathcal{F}_1 \subset \mathcal{F}_2$, then $\{F^\Delta : F \in \mathcal{F}_1\} \subseteq \{F^\Delta : F \in \mathcal{F}_2\}$ whence $\Phi(\mathcal{F}_1) \subseteq \Phi(\mathcal{F}_2)$. On the other hand, if $\mathcal{U}_1, \mathcal{U}_2 \in \mathcal{DU}(L)$ and $\mathcal{U}_1 \subset \mathcal{U}_2$, then the topology induced by $\mathcal{U}_1$ is coarser than the one induced by $\mathcal{U}_2$, hence $\Psi(\mathcal{U}_1) \subseteq \Psi(\mathcal{U}_2)$.

Finally we show that $\Phi = \Psi^{-1}$, so that Ψ is one-to-one. Given $\mathcal{F} \in \mathcal{FND}(L)$, we consider any $\mathcal{U} \in \mathcal{DU}(L)$ such that $\mathcal{F} = \Psi(\mathcal{U})$ and prove that $\Phi(\mathcal{F}) = \mathcal{U}$. If $F \in \mathcal{F}$, then it is a neighbourhood of 0, hence there is $U \in \mathcal{U}$ such that $U(0) \subseteq F$. By uniform continuity of Δ , there exists $V \in \mathcal{U}$ with $V \Delta \Delta \subset U$. Now let $(a,b) \in V$. We have $(0, a \Delta b) = (a \Delta a, b \Delta a) \in V \Delta \Delta \subset U$, whence $a \Delta b \in U(0) \subseteq F$. Hence $V \subset F^\Delta$ and therefore $\mathcal{U}$ is finer than $\Phi(\mathcal{F})$. Conversely let $U \in \mathcal{U}$. Consider a symmetric $V_1 \in \mathcal{U}$ with $V_1 \circ V_1 \subset U$, and take $V_2, V_3 \in \mathcal{U}$ such that $V_2 \vee \Delta \subset V_1$ and $V_3 \oplus \Delta \subset V_2$. Put $F = V_3(0)$, so that $F \in \mathcal{F}$. If $(a,b) \in F^\Delta$, we have $a \Delta b \in F$, that is $(0, a \Delta b) \in V_3$. It follows that $(a \wedge b, a \vee b) = (0 \oplus (a \wedge b), (a \Delta b) \oplus (a \wedge b)) \in V_3 \oplus \Delta \subset V_2$, hence $(a, a \vee b) = ((a \wedge b) \vee a, (a \vee b) \vee a) \in V_2 \vee \Delta \subset V_1$ and, similarly $(b, a \vee b) \in V_1$. Since $V_1^{-1} = V_1$ we also have $(a \vee b, b) \in V_1$, and then $(a,b) \in V_1 \circ V_1 \subset U$. Therefore $F^\Delta \subseteq U$. We conclude that $\mathcal{U} \subset \Phi(\mathcal{F})$, whence the equality. $\square$

The reader should note that the above theorem implies, as particular cases, the results of [6, Theor. 2.1] and [13, Theor. 3.6] for MV-algebras, as well as [16, Theor. 1.1] for orthomodular lattices.

From Theorem 2.4(c), by restricting to principal filters, one can deduce the order isomorphism between D-congruences and D-ideals, which has been found, using a different approach, in [4, Theor. 4.5].

2.5. Proposition. *Let $\mathcal{F}$ be the filter of neighbourhoods of 0 in a D-uniformity $\mathcal{U}$. For every $F \in \mathcal{F}$, let $F^\oplus = \{(a,b) \in L \times L : \exists h, k \in F : h \perp a, k \perp b, a \oplus h = b \oplus k\}$ and $F^\ominus = \{(a,b) \in L \times L : \exists i, j \in F : i \leq a, j \leq b, a \ominus i = b \ominus j\}$. Then both $\{F^\oplus : F \in \mathcal{F}\}$ and $\{F^\ominus : F \in \mathcal{F}\}$ are bases for $\mathcal{U}$.*

PROOF. It suffices to show that, for every $F \in \mathcal{F}$, there exist $F_1, F_2 \in \mathcal{F}$ such that $F^\oplus, F^\ominus \supseteq F_1^\Delta$ and $F_2^\oplus, F_2^\ominus \subseteq F^\Delta$.

Let $F_1 \in \mathcal{F}$ satisfy 2.3(i). Given $(a, b) \in F_1^\Delta$, we put $h = (a \vee b) \ominus a$, $k = (a \vee b) \ominus b$, $i = a \ominus (a \wedge b)$ and $j = b \ominus (a \wedge b)$. Since $h \leq (a \vee b) \ominus (a \wedge b) = a \Delta b \in F_1$, we have $h \in F$. In the same way one sees that k, i and j belong to F , too. Moreover we have $a \oplus h = a \oplus ((a \vee b) \ominus a) = a \vee b = b \oplus ((a \vee b) \ominus b) = b \oplus k$, so that $(a, b) \in F^\oplus$. Similarly, applying 1.1(i), we have $a \ominus i = a \ominus (a \ominus (a \wedge b)) = a \wedge b = b \ominus (b \ominus (a \wedge b)) = b \ominus j$, so that $(a, b) \in F^\ominus$.

Now let $G \in \mathcal{F}$ satisfy 2.3(ii), and take $F_2 \in \mathcal{F}$ satisfying 2.3(i) with G in place of F . Given $(a, b) \in F_2^\oplus$, there are $h, k \in F_2$ such that $h \perp a$, $k \perp b$ and $a \oplus h = b \oplus k$. Since $a \vee b \leq (a \oplus h) \vee (b \oplus k) = a \oplus h = b \oplus k$, we get $(a \vee b) \ominus a \leq h$ and $(a \vee b) \ominus b \leq k$, so that both $(a \vee b) \ominus a$ and $(a \vee b) \ominus b$ belong to G . By 1.1(viii), we have $a \Delta b = ((a \vee b) \ominus a) \vee ((a \vee b) \ominus b)$ hence $a \Delta b \in F$, i.e. $(a, b) \in F^\Delta$. Similarly, given $(a, b) \in F_2^\ominus$, take $i, j \in F_2$ such that $i \leq a$, $j \leq b$ and $a \ominus i = b \ominus j$. Observe that $a \ominus i = (a \ominus i) \wedge (b \ominus j) \leq a \wedge b$ thus, applying 1.1(i), $i = a \ominus (a \ominus i) \geq a \ominus (a \wedge b)$. It follows that $a \ominus (a \wedge b) \in G$, and in the same way one sees that $b \ominus (a \wedge b) \in G$, too. By 1.1(ix), we have $a \Delta b = (a \ominus (a \wedge b)) \vee (b \ominus (a \wedge b))$ hence $a \Delta b \in F$, i.e. $(a, b) \in F^\Delta$. $\square$

Given $F, G \subset L$, we will put $F \oplus G = \{f \oplus g : f \perp g, f \in F, g \in G\}$. Using this notation, condition (F1) may be rewritten as follows: $\forall F \in \mathcal{F} \quad \exists F' \in \mathcal{F} : F' \oplus F' \subseteq F$.

2.6. Proposition.

- (a) If $\mathcal{F}, \mathcal{G} \in \mathcal{FND}(L)$, then $\{F \oplus G : F \in \mathcal{F}, G \in \mathcal{G}\}$ is a base for $\mathcal{F} \wedge \mathcal{G}$ in $\mathcal{FND}(L)$.
- (b) If $\Gamma \subset \mathcal{FND}(L)$, then $\bigvee \Gamma$ in $\mathcal{FND}(L)$ is the set of all intersections of finite subsets of $\bigcup \Gamma$. In particular $\mathcal{G}_1 \vee \mathcal{G}_2 = \{G_1 \cap G_2 : G_1 \in \mathcal{G}_1, G_2 \in \mathcal{G}_2\}$ for all $\mathcal{G}_1, \mathcal{G}_2 \in \mathcal{FND}(L)$.

PROOF.

- (a) First observe that
$$\forall F \in \mathcal{F} \quad \forall G \in \mathcal{G} \quad F \cup G \subset F \oplus G. \quad (2;2)$$

Indeed, since $0 \in G$, one has $F = \{f \oplus 0 : f \in F\} \subseteq \{f \oplus g : f \perp g, f \in F, g \in G\} = F \oplus G$, and similarly for G . In particular, all sets $F \oplus G$ with $F \in \mathcal{F}$ and $G \in \mathcal{G}$ are non-empty. Now, given $F_1 \oplus G_1$ and $F_2 \oplus G_2$, with $F_1, F_2 \in \mathcal{F}$ and $G_1, G_2 \in \mathcal{G}$, let $F = F_1 \cap F_2$ and $G = G_1 \cap G_2$. We have $F \oplus G = \{f \oplus g : f \perp g, f \in F, g \in G\} \subseteq \{f \oplus g : f \perp g, f \in F_1, g \in G_1\} = F_1 \oplus G_1$ and, similarly, $F \oplus G \subset F_2 \oplus G_2$. Hence $F \oplus G \subset (F_1 \oplus G_1) \cap (F_2 \oplus G_2)$. Therefore $\{F \oplus G : F \in \mathcal{F}, G \in \mathcal{G}\}$ is a base for a filter which we denote by $\mathcal{H}$.

We prove that $\mathcal{H}$ is a D-filter. Given any $H \in \mathcal{H}$, let $F \in \mathcal{F}$ and $G \in \mathcal{G}$ such that $F \oplus G \subset H$. Take $F', F'' \in \mathcal{F}$ satisfying (F1) and (F2) respectively, and choose $G', G'' \in \mathcal{G}$ in a similar way. Clearly $H' = F' \oplus G'$ and $H'' = F'' \oplus G''$ belong to $\mathcal{H}$. We show that H' satisfies (F1) and H'' satisfies (F2) (with H in place of F). If a and b are orthogonal elements of H' , then $a = f_1 \oplus g_1$ and $b = f_2 \oplus g_2$, where $f_1, f_2 \in F'$ and $g_1, g_2 \in G'$. Note that $f_1 \perp f_2$ and $g_1 \perp g_2$, hence $f = f_1 \oplus f_2 \in F$ and $g = g_1 \oplus g_2 \in G$. Therefore $a \oplus b = (f_1 \oplus g_1) \oplus (f_2 \oplus g_2) = (f_1 \oplus f_2) \oplus (g_1 \oplus g_2) = f \oplus g \in F \oplus G \subset H$. Now let $a \in H''$ and $c \in L$. Let $f \in F''$ and $g \in G''$ such that $a = f \oplus g$, and put $d = (f \vee c) \ominus f$. We have $f' = (f \vee c) \ominus c \in F$ and $g' = (g \vee d) \ominus d \in G$. Since $g \vee d = g' \oplus d$ and $f \vee c = f \oplus d$, applying 1.1(x) and 1.1(vii), we obtain $(a \vee c) \ominus c = (a \vee f \vee c) \ominus c = ((f \oplus g) \vee (f \vee c)) \ominus c = ((f \oplus g) \vee (f \oplus d)) \ominus c = (f \oplus (g \vee d)) \ominus c = (f \oplus (g' \oplus d)) \ominus c = (f \oplus d) \oplus g' \ominus c = ((f \vee c) \oplus g') \ominus c = ((f \vee c) \ominus c) \oplus g' = f' \oplus g' \in F \oplus G \subset H$.

It follows from (2;2) that both $\mathcal{F}$ and $\mathcal{G}$ are finer than $\mathcal{H}$. To complete the proof, consider any D-filter $\mathcal{H}'$ such that both $\mathcal{F}$ and $\mathcal{G}$ are finer than $\mathcal{H}'$. We show that $\mathcal{H}' \subseteq \mathcal{H}$. Let $H \in \mathcal{H}$. By (F1), there exists $H' \in \mathcal{H}'$ such that $H' \oplus H' \subseteq H$. Since $H' \in \mathcal{F} \cap \mathcal{G}$ we get $H' \oplus H' \in \mathcal{H}$ and hence $H \in \mathcal{H}$, too.

- (b) Let $\mathcal{F}$ be the set of intersections of finite subsets of $\bigcup \Gamma$. We show that $\mathcal{F}$ is a filter.

Let $F_1, F_2 \in \mathcal{F}$. One has $F_1 = \bigcap \mathcal{F}_1$ and $F_2 = \bigcap \mathcal{F}_2$, where $\mathcal{F}_1$ and $\mathcal{F}_2$ are finite subsets of $\bigcup \Gamma$. If $G = F_1 \cap F_2$, then $G \in \mathcal{F}$ because it is the intersection of $\mathcal{F}_1 \cup \mathcal{F}_2$, which is again a finite subset of $\bigcup \Gamma$. Now let $F \in \mathcal{F}$. Then $F = \bigcap_{i=1}^n F_i$, where $F_i \in \mathcal{G}_i$ and $\mathcal{G}_i \in \Gamma$ for each

$i \in \{1, 2, \dots, n\}$. If $G \supset F$, let $A = G \setminus F$. For each i , one has $G_i = A \cup F_i \in \mathcal{G}_i$, and $\bigcap_{i=1}^n G_i = \bigcap_{i=1}^n (A \cup F_i) = A \cup \bigcap_{i=1}^n F_i = A \cup F = G$. Hence $G \in \mathcal{F}$.

Now we check properties (F1) and (F2). Let $F \in \mathcal{F}$. As above, $F = \bigcap_{i=1}^n F_i$, with $F_i \in \mathcal{G}_i \in \Gamma$. For each i , take F'_i and G_i in $\mathcal{G}_i$ satisfying (F1) and (F2) respectively (with F_i in place of F). Put $F' = \bigcap_{i=1}^n F'_i$ and $G = \bigcap_{i=1}^n G_i$. Clearly F' and G belong to $\mathcal{F}$. We show that F' satisfies (F1) and G satisfies (F2). If a and b are orthogonal elements of F' , then for each $i \in \{1, 2, \dots, n\}$ we have $a, b \in F'_i$ and hence $a \oplus b \in F_i$. Therefore $a \oplus b \in F$. Similarly, if $a \in G$ and $c \in L$, then for each $i \in \{1, 2, \dots, n\}$ we have $a \in G_i$ and hence $(a \vee c) \ominus c \in F_i$. Therefore $(a \vee c) \ominus c \in F$.

Since it is clear that each $\mathcal{G} \in \Gamma$ is contained in $\mathcal{F}$ (indeed every G in $\mathcal{G}$ is the intersection of $\{G\}$, which a finite subset of $\bigcup \Gamma$), it remains to prove that any D-filter which is finer than all filters in Γ is finer than $\mathcal{F}$, too. So let $\mathcal{G}' \in \mathcal{FND}(L)$ such that $\mathcal{G} \subset \mathcal{G}'$ for every $\mathcal{G} \in \Gamma$. Given $F \in \mathcal{F}$, one has $F = \bigcap_{i=1}^n F_i$ where $F_i \in \mathcal{G}_i \in \Gamma$, hence $F_i \in \mathcal{G}'$, for each $i \in \{1, 2, \dots, n\}$. Since $\mathcal{G}'$ is a filter, we have $F \in \mathcal{G}'$. We conclude that $\mathcal{F} \subset \mathcal{G}'$. $\square$

2.7. Corollary. $\mathcal{DU}(L)$ and $\mathcal{FND}(L)$ are distributive (complete) lattices.

PROOF. By Theorem 2.4(c), it is enough to consider $\mathcal{FND}(L)$. Let $\mathcal{F}_1, \mathcal{F}_2$ and $\mathcal{G}$ be D-filters. We have to verify that $(\mathcal{F} \vee \mathcal{G}_1) \wedge (\mathcal{F} \vee \mathcal{G}_2) \subseteq \mathcal{F} \vee (\mathcal{G}_1 \wedge \mathcal{G}_2)$.

Given $H \in (\mathcal{F} \vee \mathcal{G}_1) \wedge (\mathcal{F} \vee \mathcal{G}_2)$, take $F_1, F_2 \in \mathcal{F}$ and $G_1, G_2 \in \mathcal{G}$ with $(F_1 \cap G_1) \oplus (F_2 \cap G_2) \subseteq H$. Put $F = F_1 \cap F_2$ and let $F' \in \mathcal{F}$ satisfying 2.3(i). We complete the proof by showing that $F' \cap (G_1 \oplus G_2) \subseteq (F_1 \cap G_1) \oplus (F_2 \cap G_2)$.

Let $a \in F' \cap (G_1 \oplus G_2)$. Choose $a_1 \in G_1$ and $a_2 \in G_2$ such that $a = a_1 \oplus a_2$. Since $a_1 \leq a$ and $a \in F'$, one has $a_1 \in F \subset F_1$ and hence $a_1 \in F_1 \cap G_1$. Similarly one sees that $a_2 \in F_2 \cap G_2$. Therefore $a = a_1 \oplus a_2 \in (F_1 \cap G_1) \oplus (F_2 \cap G_2)$. $\square$

2.8. Proposition. If $\mathcal{F}, \mathcal{G} \in \mathcal{FND}(L)$, then $\{F \wedge G : F \in \mathcal{F}, G \in \mathcal{G}\}$ is a base for $\mathcal{F} \vee \mathcal{G}$, where $F \wedge G = \{f \wedge g : f \in F, g \in G\}$.

PROOF. Given $F \in \mathcal{F}$ and $G \in \mathcal{G}$, since $F \cap G = \{a \wedge a : a \in F \cap G\} \subseteq \{f \wedge g : f \in F, g \in G\} = F \wedge G$, it remains to prove that there exist $F' \in \mathcal{F}$ and $G' \in \mathcal{G}$ such that $F' \wedge G' \subseteq F \cap G$. Take $F' \in \mathcal{F}$ satisfying 2.3(i), and let G' be a member of $\mathcal{G}$ satisfying 2.3(i) also, but with G in place of F . If $f \in F'$ and $g \in G'$, then $f \wedge g \leq f$ hence $f \wedge g \in F$ and, similarly, $f \wedge g \leq g$ hence $f \wedge g \in G$. Therefore $f \wedge g \in F \cap G$. $\square$

3 Generating D-uniformities by means of k -submeasures

3.1. Definition. Let $k \geq 1$. We say that a function $\eta : L \rightarrow [0, +\infty]$ is a k -submeasure if the following conditions hold:

- (S1) $\eta(0) = 0$;
- (S2) $\forall a, b \in L \quad [a \leq b \implies \eta(a) \leq \eta(b)]$;
- (S3) $\forall a, b \in L \quad [a \perp b \implies \eta(a \oplus b) \leq k\eta(a) + \eta(b)]$;
- (S4) $\forall a, b \in L \quad \eta((a \vee b) \ominus b) \leq k\eta(a)$.

A 1-submeasure is simply called a *submeasure*.

Observe that, if L is an MV-algebra, then every function $\eta : L \rightarrow [0, +\infty]$ satisfying (S1), (S2) and (S3) with $k = 1$ is a submeasure.

For every $\varepsilon > 0$, put $S_\varepsilon = \{(x, y) \in [0, +\infty[\times [0, +\infty[: |x - y| < \varepsilon\} \cup \{(+\infty, +\infty)\}$. Then $\{S_\varepsilon : \varepsilon > 0\}$ is base for a uniformity $\mathcal{S}$ on $[0, +\infty]$ whose relativization to $[0, +\infty[$ is the usual uniformity, while $+\infty$ is a uniformly isolated point. In the sequel we will endow $[0, +\infty]$ with this uniformity.

3.2. Proposition. For every k -submeasure η there exists a D-uniformity $\mathcal{U}(\eta)$ which is the weakest D-uniformity making η uniformly continuous.

PROOF. For each $\varepsilon > 0$, let $F_\varepsilon = \{a \in L : \eta(a) < \varepsilon\}$. Since $F_{\varepsilon_1} \cap F_{\varepsilon_2} = F_{\min\{\varepsilon_1, \varepsilon_2\}}$, the collection $\{F_\varepsilon : \varepsilon > 0\}$ is a base for a filter $\mathcal{F}$. We show that $\mathcal{F}$ is a D-filter. Fix F in $\mathcal{F}$, and take $\varepsilon > 0$ with $F_\varepsilon \subset F$. Then $F' = F_{\frac{\varepsilon}{k+1}}$ satisfies (F1) and $G = F_{\frac{\varepsilon}{k}}$ satisfies (F2).

From Theorem 2.4(b), the sets F_ε^Δ form a base for a D-uniformity $\mathcal{U}(\eta)$. Now we show that η is $\mathcal{U}(\eta)$ -uniformly continuous. Let $\varepsilon > 0$ and choose $\delta = \frac{\varepsilon}{k}$. For every $(a, b) \in F_\delta^\Delta$, we have $\eta(a \vee b) = \eta((a \triangle b) \oplus (a \wedge b)) \leq k\eta(a \triangle b) + \eta(a \wedge b) < \eta(a \wedge b) + k\delta = \eta(a \wedge b) + \varepsilon$. Thus, if $\eta(a \vee b) = +\infty$, then $\eta(a \wedge b) = +\infty$ whence, by monotonicity, $\eta(a) = \eta(b) = +\infty$. Otherwise, again by monotonicity, $\eta(a)$ and $\eta(b)$ are both finite, and moreover $|\eta(a) - \eta(b)| \leq |\eta(a \vee b) - \eta(a \wedge b)| < \varepsilon$. Hence, in any case, $(\eta(a), \eta(b)) \in S_\varepsilon$.

Finally, let $\mathcal{V}$ be a D-uniformity on L making η uniformly continuous. We prove that $\mathcal{U}(\eta) \leq \mathcal{V}$, which, by Theorem 2.4(c), is equivalent to $\mathcal{F} \subset \mathcal{G}$, where $\mathcal{G}$ is the filter of neighbourhoods of 0 in $\mathcal{V}$. Take any $F \in \mathcal{F}$, and choose $\varepsilon > 0$ with $F_\varepsilon \subset F$. Since η is continuous at 0 with respect to $\mathcal{V}$, and $\eta(0) = 0$, there is some $G \in \mathcal{G}$ such that if $a \in G$ then $\eta(a) < \varepsilon$, i.e. $a \in F_\varepsilon$. It follows that $G \subset F_\varepsilon \subset F$, hence $F \in \mathcal{G}$. $\square$

Our aim is to prove a sort of converse of the previous result, namely Theorem 3.4 below.

3.3. Proposition. *Let $k, m \geq 1$, and d be a pseudometric such that for all $a, b, c \in L$:*

- (P1) $d(a \wedge c, b \wedge c) \leq d(a, b)$;
- (P2) $a \perp c, b \perp c \implies d(a \oplus c, b \oplus c) \leq kd(a, b)$;
- (P3) $d((a \vee c) \ominus c, (b \vee c) \ominus c) \leq md(a, b)$;
- (P4) $d((a \vee c) \ominus c, 0) \leq kd(a, 0)$.

For each $a \in L$, put $\tilde{\eta}(a) = d(a, 0)$. Then $\tilde{\eta}$ is a k -submeasure and $\mathcal{U}(\tilde{\eta})$ coincides with the uniformity induced by d .

PROOF. It is clear that $\tilde{\eta}$ satisfies (S1). Moreover, if $a \leq b$, by (P1) we have $\tilde{\eta}(a) = d(a, 0) = d(b \wedge a, 0 \wedge a) \leq d(b, 0) = \tilde{\eta}(b)$ and (S2) is proved. Now if $a, b \in L$ are orthogonal, then, applying the triangular inequality and (P2), we get $\tilde{\eta}(a \oplus b) = d(a \oplus b, 0) \leq d(a \oplus b, b) + d(b, 0) \leq kd(a, 0) + d(b, 0) = k\tilde{\eta}(a) + \tilde{\eta}(b)$, that is (S3). Similarly, taking any $a, b \in L$, by (P4) we get $\tilde{\eta}((a \vee b) \ominus b) = d((a \vee b) \ominus b, 0) \leq kd(a, 0) = k\tilde{\eta}(a)$, that is (S4).

Denote by $\mathcal{V}$ the uniformity induced by d . The sets $V_\varepsilon = \{(a, b) \in L \times L : d(a, b) < \varepsilon\}$ form a base for $\mathcal{V}$, while the sets $F_\varepsilon^\Delta = \{(a, b) \in L \times L : \tilde{\eta}(a \triangle b) < \varepsilon\}$ form a base for $\mathcal{U}(\tilde{\eta})$, as we have seen in Proposition 3.2. We show that for every $\varepsilon > 0$ there exists $\delta > 0$ such that $F_\delta^\Delta \subseteq V_\varepsilon$ and $V_\delta \subset F_\varepsilon^\Delta$. This will prove that $\mathcal{V} = \mathcal{U}(\tilde{\eta})$.

Take $\delta = \frac{\varepsilon}{2km}$. Given $(a, b) \in F_\delta^\Delta$, applying (P1) and (P2), we have $d(a, b) \leq d(a, a \wedge b) + d(a \wedge b, b) = d((a \vee b) \wedge a, (a \wedge b) \wedge a) + d((a \wedge b) \wedge b, (a \vee b) \wedge b) \leq 2d(a \vee b, a \wedge b) = 2d((a \triangle b) \oplus (a \wedge b), 0 \oplus (a \wedge b)) \leq 2kd(a \triangle b, 0) = 2k\tilde{\eta}(a \triangle b) < 2k\delta \leq \varepsilon$, so that $(a, b) \in V_\varepsilon$. Therefore $F_\delta^\Delta \subseteq V_\varepsilon$.

Now let $(a, b) \in V_\delta$. Recall that, by 1.1(ii), $(a \triangle b) \ominus ((a \vee b) \ominus a) = a \ominus (a \wedge b)$ and, by 1.1(viii), $(a \ominus (a \wedge b)) \wedge (b \ominus (a \wedge b)) = 0$. Hence, applying first the triangle inequality and then (P2), (P3), (P1) and again (P3), we obtain $\tilde{\eta}(a \triangle b) = d(a \triangle b, 0) \leq d(a \triangle b, (a \vee b) \ominus a) + d((a \vee b) \ominus a, 0) = d(((a \vee b) \ominus a) \oplus (a \ominus (a \wedge b)), (a \vee b) \ominus a) + d((a \vee b) \ominus a, (a \vee a) \ominus a) \leq kd(a \ominus (a \wedge b), 0) + md(a, b) = kd((a \ominus (a \wedge b)) \wedge (a \ominus (a \wedge b)), (b \ominus (a \wedge b)) \wedge (a \ominus (a \wedge b))) + md(a, b) \leq kd(a \ominus (a \wedge b), b \ominus (a \wedge b)) + md(a, b) \leq kmd(a, b) + md(a, b) < (k+1)m\delta \leq \varepsilon$ so that $(a, b) \in F_\varepsilon^\Delta$. We conclude that $V_\delta \subset F_\varepsilon^\Delta$. $\square$

Recall that if $\mathbb{G}$ is a topological Abelian group, then a mapping $\mu: L \rightarrow \mathbb{G}$ is called a *modular measure* if the following hold, for all $a, b \in L$:

- (M1) $\mu(a) + \mu(b) = \mu(a \vee b) + \mu(a \wedge b)$.
- (M2) If $a \perp b$, then $\mu(a \oplus b) = \mu(a) + \mu(b)$.

Moreover, (see [1, Theor. 3.2]) the sets $\{(a, b) \in L \times L : \forall r \leq a \triangle b \quad \mu(r) \in W\}$, where W is a neighbourhood of 0 in $\mathbb{G}$, form a base for a D-uniformity $\mathcal{U}$. This $\mathcal{U}$ is called the D-uniformity generated by μ . Note that, in case μ is positive real-valued (hence in particular a submeasure), $\mathcal{U}$ agrees with the $\mathcal{U}(\mu)$ constructed in Proposition 3.2.

3.4. Theorem. *Let $\mathcal{U}$ be a D-uniformity on L . Then:*

- (a) *For every $k > 1$ there is a family $\{\tilde{\eta}_\lambda\}_{\lambda \in \Lambda}$ of k -submeasures with $\mathcal{U} = \sup_{\lambda \in \Lambda} \mathcal{U}(\tilde{\eta}_\lambda)$. Moreover, if $\mathcal{U}$ has a countable base, we can choose $|\Lambda| = 1$.*
- (b) *If $\mathcal{U}$ is generated by a modular measure $\mu : L \rightarrow \mathbb{G}$, where $\mathbb{G}$ is a topological Abelian group, then there is a family $\{\tilde{\eta}_\lambda\}_{\lambda \in \Lambda}$ of submeasures with $\mathcal{U} = \sup_{\lambda \in \Lambda} \mathcal{U}(\tilde{\eta}_\lambda)$.*
- (c) *If L is an MV-algebra, there is a family $\{\tilde{\eta}_\lambda\}_{\lambda \in \Lambda}$ of submeasures with $\mathcal{U} = \sup_{\lambda \in \Lambda} \mathcal{U}(\tilde{\eta}_\lambda)$.*

PROOF.

(a) For every $a, b \in L$, put $f(a, b) = a \wedge b$, $g(a, b) = (a \wedge b^\perp) \oplus b$ and $h(a, b) = (a \vee b) \ominus b$. By [15, Prop. 1.1(b)], $\mathcal{U}$ has base consisting of sets U such that, for every $(a, a') \in U$ and every $b \in L$, $(f(a, b), f(a', b)) = (f(b, a), f(b, a')) \in U$. Since g and h are $\mathcal{U}$ -uniformly continuous, from [15, Prop. 1.2] it follows that $\mathcal{U}$ is generated by a family $\{d_\lambda\}_{\lambda \in \Lambda}$ of pseudometrics (a single pseudometric if Λ is countable) such that, for every $\lambda \in \Lambda$ and all $a, a', b, b' \in L$:

$$\begin{aligned} d_\lambda(f(a, b), f(a', b')) &\leq d_\lambda(a, a') + d_\lambda(b, b'), \\ d_\lambda(g(a, b), g(a', b')) &\leq k(d_\lambda(a, a') + d_\lambda(b, b')), \\ d_\lambda(h(a, b), h(a', b')) &\leq k(d_\lambda(a, a') + d_\lambda(b, b')). \end{aligned}$$

Clearly each d_λ satisfies (P1) and (P2), as well as (P3) with $m = k$, hence also (P4). Therefore, applying Proposition 3.3, the conclusion follows.

(b) Let $\{p_\lambda\}_{\lambda \in \Lambda}$ be a family of group seminorms generating the topology of $\mathbb{G}$. By [12, Theor. 3], $\mathcal{U}$ is generated by the family of pseudometrics $\{d_\lambda\}_{\lambda \in \Lambda}$ where, for every $\lambda \in \Lambda$,

$$d_\lambda(a, b) = \sup\{p_\lambda(\mu(r) - \mu(s)) : r, s \in [a \wedge b, a \vee b]\}.$$

Moreover d_λ satisfies (P1) and the following:

$$\forall a, b, c \in L \quad d_\lambda(a \vee c, b \vee c) \leq d_\lambda(a, b). \quad (3; 1)$$

We can complete the proof, applying Proposition 3.3, once we have shown that each d_λ satisfies both (P2) and (P3) with $m = k = 1$, hence also (P4).

Fix $\lambda \in \Lambda$. Given $a, b \in L$, observe first that $d_\lambda(a, b) = \sup\{p_\lambda(\mu(r)) : r \leq a \triangle b\}$. Now let $c \in L$. By Lemma 2.2 we have $((a \vee c) \ominus c) \triangle ((b \vee c) \ominus c) = (a \vee c) \triangle (b \vee c)$. Therefore, by (3; 1), $d_\lambda((a \vee c) \ominus c, ((b \vee c) \ominus c)) = d_\lambda(a \vee c, b \vee c) \leq d_\lambda(a, b)$. Finally, if $c \perp a$ and $c \perp b$, then, again by 2.2, we have $(a \oplus c) \triangle (b \oplus c) = a \triangle b$. Hence $d_\lambda(a \oplus c, b \oplus c) = d_\lambda(a, b)$.

(c) Define f , g and h as in the proof of (a). By [15, Prop. 1.5], since g is associative and distributive with respect to f , the uniformity $\mathcal{U}$ has a base consisting of sets U such that, for every $(a, a') \in U$ and every $b \in L$, $(f(a, b), f(a', b)) = (f(b, a), f(b, a')) \in U$ and $(g(a, b), g(a', b)) = (g(b, a), g(b, a')) \in U$. Moreover h is $\mathcal{U}$ -uniformly continuous, and therefore from [15, Prop. 1.2] it follows that, for any $m > 1$, $\mathcal{U}$ is generated by a family $\{d_\lambda\}_{\lambda \in \Lambda}$ of pseudometrics (a single pseudometric if Λ is countable) such that, for every $\lambda \in \Lambda$ and all $a, a', b, b' \in L$:

$$\begin{aligned} d_\lambda(f(a, b), f(a', b')) &\leq d_\lambda(a, a') + d_\lambda(b, b'), \\ d_\lambda(g(a, b), g(a', b')) &\leq (d_\lambda(a, a') + d_\lambda(b, b')), \\ d_\lambda(h(a, b), h(a', b')) &\leq m(d_\lambda(a, a') + d_\lambda(b, b')). \end{aligned}$$

Clearly each d_λ satisfies (P1), (P2) with $k = 1$ and (P3). It remains to show that (P4) with $k = 1$ is satisfied, too. Let $a, c \in L$. By (P1), we have $d_\lambda((a \vee c) \ominus c, 0) = d_\lambda(a \ominus (a \wedge c), 0) = d_\lambda((a \wedge (a \ominus (a \wedge c))), 0 \wedge (a \ominus (a \wedge c))) \leq d_\lambda(a, 0)$. $\square$

The reader should note that 3.4(c) was already proved in [6, Theor. 2.5].

References

- [1] A. Avallone. Lattice uniformities on orthomodular structures. *Mathematica Slovaca*, 51:403–419, 2001.
- [2] A. Avallone, G. Barbieri, and P. Vitolo. Hahn decomposition of modular measures and applications. Preprint.
- [3] A. Avallone and A. Basile. On a Marinacci uniqueness theorem for measures. *Journal of Mathematical Analysis and Applications*, to appear.
- [4] A. Avallone and P. Vitolo. Congruences and ideals of effect algebras. *Order*, to appear.
- [5] A. Avallone and H. Weber. Lattice uniformities generated by filters. *Journal of Mathematical Analysis and Applications*, 209(2):507–528, 1997.
- [6] G. Barbieri and H. Weber. A topological approach to the study of fuzzy measures. In *Functional Analysis and Economic Theory, Samos, 1996*, pages 17–46. Springer Verlag, Berlin, 1998.
- [7] E. G. Beltrametti and G. Cassinelli. *The Logic of Quantum Mechanics*. Addison-Wesley Publ. Co., Reading, Mass., 1981.
- [8] M. K. Bennet and D. J. Foulis. Effect algebras and unsharp quantum logics. *Found. Phys.*, 24(10):1331–1352, 1994.
- [9] D. Butnariu and P. Klement. *Triangular Norm-based Measures and Games with Fuzzy Coalitions*. Kluwer Academic Publishers, Dordrecht, 1993.
- [10] A. Dvurečenskij and S. Pulmannová. *New Trends in Quantum Structures*. Kluwer Academic Publishers, Dordrecht, 2000.
- [11] L. G. Epstein and J. Zhang. Subjective probabilities on subjectively unambiguous events. *Econometrica*, 69(2):265–306, 2001.
- [12] I. Fleischer and T. Traynor. Group-valued modular functions. *Algebra Universalis*, 14:287–291, 1982.
- [13] M. G. Graziano. Uniformities of Fréchet-Nikodym type on Vitali spaces. *Semigroup Forum*, 61(1):91–115, 2000.
- [14] H. Weber. Uniform lattices. I: A generalization of topological Riesz spaces and topological Boolean rings. *Annali di Matematica Pura e Applicata (4)*, 160:347–370, 1991.
- [15] H. Weber. Metrization of uniform lattices. *Czechoslovak Mathematical Journal*, 118:271–280, 1993.
- [16] H. Weber. Lattice uniformities and modular functions on orthomodular lattices. *Order*, 12:295–305, 1995.